

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 992 510 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
12.03.2003 Bulletin 2003/11

(51) Int Cl.7: **C07H 19/167, A61K 31/70**

(21) Application number: **99119389.7**

(22) Date of filing: **20.08.1997**

(54) **N6 heterocyclic substituted adenosine derivatives**

N6-heterocyclisch substituierte Adenosin-Derivate

Dérivés d'adénosine à substitution hétérocyclique N6

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**
Designated Extension States:
AL LT LV RO SI

(30) Priority: **27.08.1996 US 702234**

(43) Date of publication of application:
12.04.2000 Bulletin 2000/15

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
97938502.8 / 0 920 438

(73) Proprietor: **CV THERAPEUTICS, INC.**
Palo Alto, CA 94304 (US)

(72) Inventors:
• **Lum, Robert T.**
Palo Alto, CA 94306 (US)
• **Pfister, Jurg R.**
Los Altos, CA 94024 (US)
• **Schow, Steven R.**
Redwood City, CA 94065 (US)
• **Wick, Michael M.**
Chestnut Hill, MA 02146 (US)
• **Nelson, Marek G.**
Sunol, CA 95051 (US)
• **Schreiner, George F.**
Los Altos Hills, CA 94022 (US)

(74) Representative: **Vossius, Volker, Dr. et al**
Dr. Volker Vossius,
Patentanwaltskanzlei - Rechtsanwaltskanzlei,
Geibelstrasse 6
81679 München (DE)

(56) References cited:
EP-A- 0 355 825 EP-A- 0 704 215
WO-A-95/11904 WO-A-97/24363

- **KNUTSEN L J S ET AL: "The synthesis and biochemical evaluation of new A1 selective adenosine receptor agonists containing 6-hydrazinopurine moieties" BIOORG. MED. CHEM. LETT. (BMCLE8,0960894X);1993; VOL.3 (12); PP.2661-6, XP000870169 Novo Nord. Pharm. Div.;Dep. Med. Chem.; Maaloev; DK 2760; Den. (DK)**
- **JACOBSON K A ET AL: "Molecular probes for extracellular adenosine receptors" BIOCHEM. PHARMACOL. (BCPCA6,00062952);1987; VOL.36 (10); PP.1697-707, XP000870192 Natl. Inst. Diabetes Dig. Kidney Dis.;Lab. Chem.; Bethesda; 20892; MD; USA (US)**
- **DALY J W ET AL: "Structure-activity relationships for N6-substituted adenosines at a brain A1-adenosine receptor with a comparison to an A2-adenosine receptor regulating coronary blood flow" BIOCHEM. PHARMACOL. (BCPCA6,00062952);1986; VOL.35 (15); PP.2467-81, XP000870166 Natl. Inst. Arthritis, Diabetes, Dig. Kidney Dis.;Lab. Bioorg. Chem.; Bethesda; 20205; MD; USA (US)**
- **MUNTER T ET AL: "Identification of Adducts Formed in Reaction of Adenosine with 3-Chloro-4-methyl-5-hydroxy-2(5H)-furanone , a Bacterial Mutagen Present in Chlorine Disinfected Drinking Water" CHEM. RES. TOXICOL. (CRTOEC,0893228X);1996; VOL.9 (4); PP.703-708, XP000870168 Aabo Akademi University;Department of Organic Chemistry; Turku; FIN-20500; Finland (FI)**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 992 510 B1

- BJOERKMAN P O ET AL: "Acetylation of cytokinins and modified adenine compounds: a simple and non-destructive derivatization method for gas chromatography-mass spectrometric analysis" PHYTOCHEM. ANAL. (PHANEL,09580344);1996; VOL.7 (2); PP.57-68, XP002128803 Swedish University Agricultural Sciences;Dep. Plant Physiology; Uppsala; S-750 07; Swed. (SE)

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Description

BACKGROUND OF THE INVENTION**Field of Invention**

[0001] This invention encompasses optionally substituted O-heterocyclic adenosine derivatives that are selective adenosine type 1 receptor agonists, and as such, are potentially useful agents for the treatment of cardiovascular diseases and central nervous system disorders.

Description of the Art

[0002] There are two subtypes of adenosine receptors in the heart: A_1 and A_2 . Each subtype effects different physiological functions. Stimulation of the A_1 adenosine receptor induces two distinct physiological responses. The first is the inhibition of the stimulatory effects of catecholamine. This effect is mediated via the inhibition of cyclic AMP synthesis. The second effect mediated by A_1 receptors is the slowing of the heart rate and impulse propagation through the AV node. The effect is independent of cAMP metabolism and is associated with A_1 adenosine receptor activation of the inwardly rectifying K^+ channel. This effect is unique to the A_1 receptor; there is no role for the A_2 receptor in modulating the function of this channel. Stimulation of the adenosine A_1 receptor accordingly shortens the duration and decreases the amplitude of the action potential of AV nodal cells and subsequently prolongs the refractory period of the cells. The consequence of these effects is to limit the number of impulses conducted from the atria to the ventricles. This forms the basis of the clinical utility of A_1 receptor agonists for the treatment of supraventricular tachycardias, including atrial fibrillation, atrial flutter, and AV nodal re-entrant tachycardia.

[0003] The clinical utility of A_1 agonists therefore would be in the treatment of acute and chronic disorders of heart rhythm, especially those diseases characterized by rapid heart rate where the rate is driven by abnormalities in the atria. The disorders include but are not limited to atrial fibrillation, supra ventricular tachycardia and atrial flutter. Exposure to A_1 agonists causes a reduction in the heart rate and a regularization of the abnormal rhythm thereby restoring improved hemodynamic blood flow.

[0004] A_1 agonists, through their ability to inhibit the catecholamine induced increase in cAMP, should have beneficial effects in the failing heart where increased sympathetic tone causing enhanced cAMP has been associated with increased likelihood of ventricular arrhythmias and sudden death.


SUMMARY OF THE INVENTION

[0005] An object of this invention is novel heterocyclic substituted adenosine derivatives.

[0006] Another object of this invention is novel heterocyclic substituted adenosine derivatives that are useful as A_1 receptor agonists.

[0007] Still another object of this invention is novel heterocyclic substituted adenosine derivatives that are useful for treating supraventricular tachycardias, including atrial fibrillation, atrial flutter, and AV nodal re-entrant tachycardia.

[0008] In one embodiment, this invention relates to a compound having the formula:



wherein R_1 is an unsubstituted or substituted monocyclic or polycyclic O-heterocyclic group containing from 3 to 15 atoms, and wherein R_1 may be mono or polysubstituted with one or more compounds selected from the group consisting of halogen, oxo, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, cyano and mixtures thereof
 5 wherein R_1 does not contain an epoxide group (see WO-A-9724363) and with the proviso that the compound is not 5-N⁶-Adenosyl-3-chloro-4-methyl-2(5H)-furanone (see Chem. Res. Toxicol; 1996, 9,703-708).

[0009] In another embodiment, this invention discloses a method for stimulating coronary activity in a mammal experiencing a cardiac electrical disorder that can be treated by stimulating an A_1 heart adenosine receptor by administering a therapeutically effective amount of the composition disclosed above to the mammal.

10 [0010] In still another embodiment, this invention relates to a pharmaceutical composition comprising the above compound and one or more pharmaceutical excipients.

DESCRIPTION OF THE FIGURES


15 [0011]

Figure 1 is a plot of the effect of the concentration compound II of Example 2 on atrial AV nodal conductance for the A_1 adenosine receptor (-●-) and for the A_2 adenosine receptor (-○-).

20 Figure 2 is a plot of the effect of the concentration of compound I of Example 2 on atrial AV nodal conductance and specifically on the response of the A_1 adenosine receptor (-●-) and on the response of the A_2 adenosine receptor (-○-).

DESCRIPTION OF THE CURRENT EMBODIMENT

25 [0012] This invention comprises adenosine derivatives which are selective adenosine type 1 receptor agonists of the formula:




45 where:

R_1 is a cycloalkyl group which is an unsubstituted or substituted monocyclic or polycyclic O-heterocyclic group with 3 to 15 atoms in the ring structure. R_1 may optionally be mono or polysubstituted with halogen, oxo, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, and cyano. However, R_1 cannot contain an epoxy group, and the compounds is not 5-N⁶-Adenosyl-3-chloro-4-methyl-2(5H)-furanone.


50 R_1 is preferably a monocyclic, bicyclic, or tricyclic group containing from 3 to 15 atoms, at least one of which is O, wherein R_1 may be mono or polysubstituted with one or more compounds selected from the group consisting of halogen, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, cyano and mixtures thereof.

55 [0013] In a more preferred embodiment, R_1 is:



wherein R_1' , R_1'' , R_1''' , and R_1'''' are individually selected from the group halogen, oxo, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, cyano and mixtures thereof. Preferably, R_1' , R_1'' , R_1''' , and R_1'''' are individually selected from the group H, lower alkyl, substituted lower alkyl, alkoxy, aryl, and substituted aryl. By "individually selected" it is meant that R_1' , R_1'' , R_1''' , and R_1'''' may each be a different component, each may be the same component, e.g., hydrogen, or some of the components may be the same and some different. It is most preferred that when R_1 is the composition set forth above, that R_1' , R_1'' , R_1''' , and R_1'''' are individually selected from the group H, lower alkyl, and substituted lower alkyl. R_1''' and R_1'''' may also be a single oxygen atom.

[0014] In an alternative embodiment, R_1 is selected from the group consisting of:



wherein each R may individually selected from the group consisting of H, lower alkyl, and substituted lower alkyl. In a most preferred embodiment, R_1 is selected from the group consisting of 3-tetrahydrofuranyl, 3-tetrahydrothiofuranyl, 4-pyranyl and 4-thiopyranyl.

[0015] The following definitions apply to terms as used herein.

[0016] The term "halogen" refers to fluorine, bromine, chlorine, and iodine atoms.

[0017] The term "oxo" refers to $=O$.

[0018] The term "hydroxyl" refers to the group $-OH$.

The term "lower alkyl" refers to a cyclic, branched or straight chain, alkyl group of one to ten carbon atoms. This term is further exemplified by such groups as methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, i-butyl (or 2-methylpropyl), cyclopropylmethyl, i-amyl, n-amyl, hexyl and the like.

[0019] The term "substituted lower alkyl" refers to lower alkyl as just described including one or more groups such as hydroxyl, thiol, alkylthiol, halogen, alkoxy, amino, amido, carboxyl, cycloalkyl, substituted cycloalkyl, heterocycle, cycloheteroalkyl, substituted cycloheteroalkyl, acyl, carboxyl, aryl, substituted aryl, aryloxy, hetaryl, substituted hetaryl, aralkyl, heteroaralkyl, alkyl alkenyl, alkyl alkynyl, alkyl cycloalkyl, alkyl cycloheteroalkyl, and cyano. These groups may be attached to any carbon atom of the lower alkyl moiety.

[0020] The term "alkoxy" refers to the group $-OR$, where R is lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroalkyl, heteroarylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, or substituted cycloheteroalkyl as defined below.

[0021] The term "acyl" denotes groups $-C(O)R$, where R is hydrogen, lower alkyl substituted lower alkyl, aryl, substituted aryl, amino, and the like as defined below.

[0022] The term "aryloxy" denotes groups $-OAr$, where Ar is an aryl, substituted aryl, heteroaryl, or substituted heteroaryl group as defined below.

[0023] The term "amino" refers to the group NR_2R_2' , where R_2 and R_2' may independently be hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, hetaryl, or substituted hetaryl as defined herein.

[0024] The term "carboxyl" denotes the group -C(O)OR, where R may independently be hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, hetaryl, substituted hetaryl and the like as defined herein.

[0025] The term "aryl" or "Ar" refers to an aromatic carbocyclic group having at least one aromatic ring (e.g., phenyl or biphenyl) or multiple condensed rings in which at least one ring is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl, anthryl, or phenanthryl).

[0026] The term "substituted aryl" refers to aryl optionally substituted with one or more functional groups, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol sulfamido and the like.

[0027] The term "heterocycle", as a substituent of R₁, refers to a saturated, unsaturated or aromatic carbocyclic group having a single ring (e.g., morpholino, pyridyl or furyl) or multiple condensed rings (e.g., naphthpyridyl, quinoxalyl, quinolinyl, indolizinyll or benzo[b]thienyl) and having at least one hetero atom, such as N, O or S, within the ring, which can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol or sulfamido.

[0028] The term "heteroaryl" or "hetar" refers to a heterocycle in which at least one heterocyclic ring is aromatic.

[0029] The term "substituted heteroaryl" refers to a heterocycle optionally mono or poly substituted with one or more functional groups, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido.

[0030] The term "cycloalkyl" refers to a divalent cyclic or polycyclic alkyl group containing 3 to 15 carbon atoms.

[0031] The term "substituted cycloalkyl" refers to a cycloalkyl group comprising one or more substituents with, e.g., halogen, lower alkyl, substituted lower alkyl, alkoxy, alkylthio, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido.

[0032] The compositions of this invention are useful as A₁ receptor agonists for the treatment of cardiac electrical disorders such as supraventricular tachycardias, including atrial fibrillation, atrial flutter, and AV nodal re-entrant tachycardia. The compositions may be administered orally, intravenously, through the epidermis or by any other means known in the art for administering a therapeutic agents.

[0033] The method of treatment comprises the administration of an effective quantity of the chosen compound, preferably dispersed in a pharmaceutical carrier. Dosage units of the active ingredient are generally selected from the range of 0.01 to 100 mg/kg, but will be readily determined by one skilled in the art depending upon the route of administration, age and condition of the patient. These dosage units may be administered one to ten times daily for acute or chronic disorders. No unacceptable toxicological effects are expected when compounds of the invention are administered in accordance with the present invention.

[0034] If the final compound of this invention contains a basic group, an acid addition salt may be prepared. Acid addition salts of the compounds are prepared in a standard manner in a suitable solvent from the parent compound and an excess of acid, such as hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, maleic, succinic, or methane sulfonic. The hydrochloric salt form is especially useful. If the final compound contains an acidic group, cationic salts may be prepared. Typically the parent compound is treated with an excess of an alkaline reagent, such as hydroxide, carbonate or alkoxide, containing the appropriate cation. Cations such as Na⁺, K⁺, Ca²⁺ and NH₄⁺ are examples of cations present in pharmaceutically acceptable salts. Certain of the compounds form inner salts or Zwitterions which may also be acceptable.


[0035] Pharmaceutical compositions including the compounds of this invention, and/or derivatives thereof, may be formulated as solutions or lyophilized powders for parenteral administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. If used in liquid form the compositions of this invention are preferably incorporated into a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water and buffered sodium or ammonium acetate solution. Such liquid formulations are suitable for parenteral administration, but may also be used for oral administration. It may be desirable to add excipients such as polyvinylpyrrolidinone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride, sodium citrate or any other excipient known to one of skill in the art to pharmaceutical compositions including compounds of this invention. Alternatively, the pharmaceutical compounds may be encapsulated, tableted or prepared in an emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Liquid carriers include syrup, peanut oil, olive oil, glycerin, saline, alcohols and water. Solid carriers include starch, lactose, calcium sulfate, dihydrate, teffa alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. The carrier may also include a sustained release material such as glycerol monostearate or glycerol distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 1 gram per dosage unit. The pharmaceutical dosages are made using conventional techniques such as milling, mixing, granulation, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous

suspension. Such a liquid formulation may be administered directly or filled into a soft gelatin capsule.

[0036] The Examples which follow serve to illustrate this invention. The Examples are intended to in no way limit the scope of this invention, but are provided to show how to make and use the compounds of this invention. In the Examples, all temperatures are in degrees Celsius.

EXAMPLE 1

[0037] The compounds of this invention may be prepared by conventional methods of organic chemistry. The reaction sequence outlined below, is a general method, useful for the preparation of compounds of this invention.



According to this method, oxacycloalkyl carboxylic acid is heated in a mixture of dioxane, diphenylphosphoryl azide and triethylamine for 1 hour. To this mixture is added benzyl alcohol and the reaction is further heated over night to give intermediate compound 1. Compound 1 is dissolved in methanol. Next, concentrated HCl, Pd/C is added and the mixture is placed under hydrogen at 1 atm. The mixture is stirred overnight at room temperature and filtered. The residue is recrystallized to give intermediate compound 2. 6-chloropurine riboside is combined and the mixture is compound 2 dissolved in methanol and treated with triethylamine. The reaction is heated to 80° C for 30 hours. Isolation and purification leads to Compound 3.

EXAMPLE 2

[0038] Compounds of this invention prepared according to the method of Example 1 were tested in two functional models specific for adenosine A₁ receptor agonist function. The first was the A₁ receptor mediated inhibition of isoproterenol stimulated cAMP accumulation in DDT cells. The EC₅₀ of each derivative is shown in Table I. Also shown in Table I is the ability of each derivative to stimulate cAMP production in PC12 cells, a function of agonist stimulation of adenosine A₂ receptors. The ratio of the relative potency of each compound in stimulating either an A₁ receptor or an A₂ receptor effect is termed the selectivity of each compound for the A₁ receptor. As can be seen in Table I, each derivative is relatively selective as an A₁ receptor agonist. The use of measuring cAMP metabolism as an assay for adenosine A₁ receptor function has been previously described (Scammells, P., Baker, S., Belardinelli, L., and Olsson,

R. , 1994, Substituted 1,3-dipropylxanthines as irreversible antagonists of A₁ adenosine receptors. J. Med. Chem 37: 2794-2712, 1994).

Table I

Compound	R	EC ₅₀ (nM) DDT cells	EC ₅₀ (nM) PC12 cells	A ₁ /A ₂	A ₂ /A ₁
I	4-aminopyran	12	970	0.012	80.0
II	(±)-3-aminotetrahydrofuran	13	1400	0.0093	107.6
III	(R)-3-aminotetrahydrofuran	1.08	448	0.0024	414
IV	(1)-caprolactam	161	181	0.889	1.12
V	(S)-3-aminotetrahydrofuran	3.40	7680	0.00044	2258

[0039] Compounds were also tested in a whole organ model of A₁ receptor activation with respect to atrial and AV nodal function. In this model, guinea pig hearts are isolated and perfused with saline containing compound while atrial rate and AV nodal conduction time are assessed by electrographic measurement of atrial cycle length and AV intervals, as detailed in Belardinelli, L, Lu, J. Dennis, D. Martens, J, and Shryock J. (1994); The cardiac effects of a novel A₁-adenosine receptor agonist in guinea pig isolated heart. J. Pharm. Exp. Therap. 271:1371-1382 (1994). As shown in Figure 1, each derivative was effective in slowing the atrial rate and prolonging the AV nodal conduction time of spontaneously beating hearts in a concentration-dependent manner, demonstrating efficacy as adenosine A₁ receptor agonists in the intact heart.

EXAMPLE 3

Preparation of N-benzyloxycarbonyl-4-aminopyran.

[0040] A mixture of 4-pyranilcarboxylic acid (2.28 gm, 20 mmol), diphenylphosphorylazide (4.31 ml, 20 mmol), triethylamine (2.78 ml, 20 mmol) in dioxane (40 ml) was heated in a 100° C oil bath under dry nitrogen for 1 hour. Benzyl alcohol (2.7 ml, 26 mmol) was added, and heating was continued at 100° C for 22 hours. The mixture was cooled, filtered from a white precipitate and concentrated. The residue was dissolved in 2N HCl and extracted twice with EtOAc. The extracts were washed with water, sodium bicarbonate, brine and then dried over MgSO₄, and concentrated to an oil which solidified upon standing. The oil was chromatographed (30% to 60% EtOAc/Hex) to give 1.85 g of a white solid (40%).

Preparation of 4-aminopyran.

[0041] N-benzyloxycarbonyl-4-aminopyran (1.85 gm, 7.87 mmol) was dissolved in MeOH (50 ml) along with conc. HCl and Pd-C (10%, 300 mg). The vessel was charged with hydrogen at 1 atm and the mixture was allowed to stir for 18 hours at room temperature. The mixture was filtered through a pad of celite and concentrated. The residue was co-evaporated twice with MeOH/EtOAc and recrystallized from MeOH/EtOAc to afford 980 mg (91 %) of white needles (mp 228-230° C).

Preparation of 6-(4-aminopyran)-purine riboside.

[0042] A mixture of 6-chloropurine riboside (0.318 gm, 1.1 mmol), 4-aminopyran-HCl (0.220 mg, 1.6 mmol) and triethylamine (0.385 ml, 2.5 mmol) in methanol (10 ml) was heated to 80° C for 30 hours. The mixture was cooled, concentrated and the residue chromatographed (90:10:1, CH₂Cl₂/MeOH/PrNH₂).

[0043] The appropriate fractions were collected and rechromatographed using a chromatotron (2 mm plate, 90: 10: 1, CH₂ Cl₂/MeOH/PrNH₂) to give an off white foam (0.37 gm, 95%).

EXAMPLE 4

Preparation of N-benzyloxycarbonyl-3-aminotetrahydrofuran.

[0044] A mixture of 3-tetrahydrofuroic acid (3.5 gm, 30 mmol), diphenylphosphorylazide (6.82 ml, 32 mmol), triethylamine (5 ml, 36 mmol) in dioxane (35 ml) was stirred at RT for 20 min then heated in a 100° C oil bath under dry

nitrogen for 2 hours. Benzyl alcohol (4.7 ml, 45 mmol) was added, and continued heating at 100° C for 22 hours. The mixture was cooled, filtered from a white precipitate and concentrated. The residue was dissolved in 2N HCl and extracted twice using EtOAc. The extracts were washed with water, sodium bicarbonate, brine dried over MgSO₄, and then concentrated to an oil which solidifies upon standing. The oil was chromatographed (30% to 60% EtOAc/Hex) to give 3.4 g of an oil (51 %).

Preparation of 3-aminotetrahydrofuran.

[0045] N-benzylloxycarbonyl-3-aminotetrahydrofuran (3.4 gm, 15 mmol) was dissolved in MeOH (50 ml) along with conc. HCl and Pd-C (10%, 300 mg). The vessel was charged with hydrogen at 1 atm and the mixture was allowed to stir for 18 hours at room temperature. The mixture was filtered through a pad of celite and concentrated. The residue was co-evaporated two times with MeOH/EtOAc and recrystallized from MeOH/EtOAc to give 1.9 g of a yellow solid.

Preparation of 6-(3-aminotetrahydrofuranyl)purine riboside.

[0046] A mixture of 6-chloropurine riboside (0.5 gm, 1.74 mmol), 3-aminotetrahydrofuran (0.325 gm, 2.6 mmol) and triethylamine (0.73 ml, 5.22 mmol) in methanol (10 ml) was heated to 80° C for 40 hours. The mixture was cooled, and concentrated. The residue was filtered through a short column of silica gel eluting with 90/10/1 (CH₂Cl₂/MeOH/PrNH₂), the fractions containing the product were combined and concentrated. The residue was chromatographed on the chromatotron (2 mm plate, 92.5/7.5/1, CH₂Cl₂/MeOH/PrNH₂). The resulting white solid was recrystallized from MeOH/EtOAc to give 0.27 gm of white crystals (mp 128-130° C).

EXAMPLE 5

Resolution of 3-aminotetrahydrofuran hydrochloride

[0047] A mixture of 3-aminotetrahydrofuran hydrochloride (0.5 gm, 4 mmol) and (S)-(+)-10-camphorsulfonyl chloride (1.1 gm, 4.4 mmol) in pyridine (10 ml) was stirred for 4 hours at room temperature and then concentrated. The residue was dissolved in EtOAc and washed with 0.5N HCl, sodium bicarbonate and brine. The organic layer was dried over MgSO₄, filtered and concentrated to give 1.17 g of a brown oil (97%) which was chromatographed on silica gel (25% to 70% EtOAc/Hex). The white solid obtained was repeatedly recrystallized from acetone and the crystals and supernatant pooled until an enhancement of greater than 90% by ¹H NMR was achieved.

Preparation of 3-(S)-aminotetrahydrofuran hydrochloride.


[0048] The sulfonamide (170 mg, 0.56 mmol) was dissolved in conc. HCl/AcOH (2 mL each), stirred for 20 hours at room temperature, washed three times with CH₂Cl₂ (10 ml) and concentrated to dryness to give 75 mg (quant.) of a white solid.

Preparation of 6-(3-(S)-aminotetrahydrofuranyl)purine riboside.

[0049] A mixture of 6-chloropurine riboside (30 mg, 0.10 mmol), 3-(S)-aminotetrahydrofuran hydrochloride (19 mg, 0.15 mmol) and triethylamine (45 ml, 0.32 mmol) in methanol (0.5 ml) was heated to 80° C for 18 hours. The mixture was cooled, concentrated and chromatographed with 95/5 (CH₂Cl₂/MeOH) to give 8 mg (24%) of a white solid.


Claims

1. A compound with a general formula



wherein R_1 is an unsubstituted or substituted monocyclic or polycyclic O-heterocyclic group with 3 to 15 atoms in the ring structure, wherein R_1 does not contain an epoxide group and with the proviso that the compound is not 5-N⁶-Adenosyl-3-chloro-4-methyl-2(SH)-furanone.

2. The compound of claim 1 wherein R_1 is mono or polysubstituted with one or more substituents selected from the group consisting of halogen, oxo, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, cyano and mixtures thereof, whereby lower alkyl refers to a cyclic, branched or straight chain alkyl group of one to ten carbon atoms.
3. The compounds N-(3(R)-tetrahydrofuranyl)-6-amino-purine riboside, N-(3(S)-tetrahydrofuranyl)-6-amino-purine riboside and N-(3-tetrahydrofuranyl)-6-amino-purine riboside.
4. The compound of claim 1 wherein R_1 is selected from the group consisting of:



wherein each R may be individually selected from the group consisting of H; lower alkyl, and substituted lower alkyl, as defined above.

5. The compound of claim 1 wherein R_1 is selected from the group consisting of 3-tetrahydrofuranyl, and 4-pyranyl.
6. An adenosine type 1 receptor agonist comprising the compound of claim 1.
7. Use of a therapeutically effective amount of a compound according to any one claims 1 to 5 for the preparation of a pharmaceutical composition for stimulating coronary activity in a mammal experiencing a coronary electrical disorder that can be treated by stimulating an A_1 heart adenosine receptor.
8. The use of claim 7 wherein the therapeutically effective amount ranges from about 0.01 to about 100 mg/kg weight of the mammal.

9. The use of claim 7 wherein the composition is to be administered to a mammal experiencing a coronary electrical disorder selected from the group consisting of supraventricular tachycardias, atrial fibrillation, atrial flutter, and AV nodal re-entrant tachycardia.

10. The use of claim 9 wherein the mammal is a human.


11. A pharmaceutical composition comprising the compound of any of claims 1 to 7 and one or more pharmaceutical excipients.

12. The pharmaceutical composition of matter of claim 11 wherein the pharmaceutical composition is in the form of a solution.

13. The pharmaceutical composition of matter of claim 11 wherein the pharmaceutical composition is in the form of a tablet.

Patentansprüche

1. Verbindung mit der allgemeinen Formel




worin R_1 eine unsubstituierte oder substituierte monocyclische oder polycyclische O-heterocyclische Gruppe mit 3 bis 15 Atomen in der Ringstruktur ist, worin R_1 keine Epoxidgruppe enthält, und mit der Maßgabe, dass die Verbindung nicht 5-N⁶-Adenosyl-3-chlor-4-methyl-2(5H)-furanon ist.

2. Verbindung nach Anspruch 1, worin R_1 mit einem oder mehreren Substituenten mono- oder polysubstituiert ist, die ausgewählt sind aus der Gruppe bestehend aus Halogenatom, Oxo-, Hydroxyl-, niederer Alkyl-, substituierter niederer Alkyl-, Alkoxy-, Aryl-, Acyl-, Aryloxy-, Carboxyl-, substituierter Aryl-, heterocyclischer, Heteroaryl-, substituierter Heteroaryl-, Cycloalkyl-, substituierter Cycloalkyl-, Nitro-, Cyanogruppe und Gemischen davon, wobei eine niedere Alkylgruppe eine cyclische, verzweigte oder unverzweigte Alkylgruppe mit 1 bis 10 Kohlenstoffatomen betrifft.

3. Die Verbindungen N-(3(R)-Tetrahydrofuranyl)-6-amino-purinribosid, N-(3(S)-Tetrahydrofuranyl)-6-amino-purinribosid und N-(3-Tetrahydrofuranyl)-6-amino-purinribosid.

4. Verbindung nach Anspruch 1, worin R_1 ausgewählt ist aus der Gruppe bestehend aus



15 worin jedes R unabhängig ausgewählt ist aus der Gruppe bestehend aus H-Atom, niederer Alkyl- und substituierter niederer Alkylgruppe, wie vorstehend definiert.

5. Verbindung nach Anspruch 1, worin R₁ ausgewählt ist aus der Gruppe bestehend aus 3-Tetrahydrofuranyl- und 4-Pyranylgruppe.

20 6. Adenosin Typ 1-Rezeptoragonist, umfassend die Verbindung nach Anspruch 1.

7. Verwendung einer therapeutisch wirksamen Menge einer Verbindung nach einem der Ansprüche 1 bis 5 für die Herstellung einer pharmazeutischen Zusammensetzung zum Stimulieren der Koronaraktivität in einem Säuger, der eine Koronarreizstörung erfährt, die durch Stimulation eines A₁-Herz-Adenosinrezeptors behandelt werden kann.

8. Verwendung nach Anspruch 7, wobei die therapeutisch wirksame Menge etwa 0,01 bis etwa 100 mg/kg Gewicht des Säugers beträgt.

9. Verwendung nach Anspruch 7, wobei die Zusammensetzung an einen Säuger verabreicht werden soll, der eine Koronarreizstörung erfährt, die ausgewählt ist aus der Gruppe bestehend aus supraventrikulären Tachykardien, Vorhofflimmern, Vorhofflattern und AV-Knoten-Reentry-Tachykardie.

10. Verwendung nach Anspruch 9, wobei der Säuger ein Mensch ist.


11. Pharmazeutische Zusammensetzung, umfassend die Verbindung nach einem der Ansprüche 1 bis 7 und ein oder mehrere pharmazeutische Exzipienzien.

12. Pharmazeutische Zusammensetzung nach Anspruch 11, wobei die pharmazeutische Zusammensetzung in Form einer Lösung vorliegt.

13. Pharmazeutische Zusammensetzung nach Anspruch 11, wobei die pharmazeutische Zusammensetzung in Form einer Tablette vorliegt.


Revendications

1. Composé ayant pour formule générale



où R_1 est un groupe O-hétérocyclique monocyclique ou polycyclique, substitué ou non-substitué, ayant 3 à 15 atomes dans la structure cyclique, tandis que R_1 ne contient pas de groupe époxyde et avec pour condition que le composé n'est pas la 5-N⁶-adénosyl-3-chloro-4-méthyl-2(5H)-furanone.

2. Composé selon la revendication 1 dans lequel R_1 est mono- ou polysubstitué par un ou plusieurs substituants choisis dans le groupe consistant en halogène, oxo, hydroxyle, alkyle inférieur, alkyle inférieur substitué, alcoxy, aryle, acyle, aryloxy, carboxyle, aryle substitué, hétérocycle, hétéroaryle, hétéroaryle substitué, cycloalkyle, cycloalkyle substitué, nitro, cyano et leurs mélanges, tandis que alkyle inférieur désigne un groupe alkyle cyclique, à chaîne droite ou à chaîne ramifiée, ayant un à dix atomes de carbone.
3. Composés N-(3(R)-tétrahydrofuranyl)-6-amino-purine riboside, N-(3(S)-tétrahydrofuranyl)-6-amino-purine riboside et N-(3-tétrahydrofuranyl)-6-amino-purine riboside.
4. Composé selon la revendication 1 dans lequel R_1 est choisi dans le groupe consistant en:



où chaque R peut être choisi individuellement dans le groupe consistant en H, alkyle inférieur et alkyle inférieur substitué, comme définis plus haut.

5. Composé selon la revendication 1 dans lequel R_1 est choisi dans le groupe consistant en 3-tétrahydrofuranyle et 4-pyranyle.
6. Agoniste du récepteur de type 1 d'adénosine comprenant le composé selon la revendication 1.
7. Utilisation d'une quantité thérapeutiquement efficace d'un composé selon l'une quelconque des revendications 1 à 5 pour la préparation d'une composition pharmaceutique pour la stimulation de l'activité coronarienne chez un

mammifère ressentant un trouble électrique coronarien qui peut être traité par stimulation d'un récepteur d'adénosine cardiaque A₁.

- 5 8. Utilisation selon la revendication 7, dans laquelle la quantité thérapeutiquement efficace se situe entre environ 0,01 et environ 100 mg/kg de poids du mammifère.
- 10 9. Utilisation selon la revendication 7, dans laquelle la composition est destinée à être administrée à un mammifère ressentant un trouble électrique coronarien choisi dans le groupe consistant en tachycardies supraventriculaires, fibrillation auriculaire, flutter auriculaire, et tachycardie ré-entrante nodale AV.
- 15 10. Utilisation selon la revendication 9, dans laquelle le mammifère est un humain.
- 20 11. Composition pharmaceutique comprenant le composé selon l'une quelconque des revendications 1 à 7 et un ou plusieurs excipients pharmaceutiques.
- 25 12. Composition pharmaceutique selon la revendication 11, dans laquelle la composition pharmaceutique est sous la forme d'une solution.
- 30 13. Composition pharmaceutique selon la revendication 11, dans laquelle la composition pharmaceutique est sous la forme d'un comprimé.
- 35
- 40
- 45
- 50
- 55

